

Search for Proton Medium Modifications in the ${}^4\text{He}(e,e'p)$ Reaction

Steffen Strauch
University of South Carolina

2007 Hall C User's Summer Workshop, August 9-10, 2007
Jefferson Lab, Newport News, VA

Outline

- Nucleon medium modifications
 - ▶ Signatures and experimental limits
 - ▶ Models for in-medium form factors
- Results from JLab $^4\text{He}(\mathbf{e},\mathbf{e}'\mathbf{p})$ experiments
 - ▶ Polarization-transfer technique
 - ▶ Competing interpretations of previous data from E93-049
 - ▶ New constraints from preliminary data* from E03-104
- Possible new experiment in Hall C
- Summary


*Simona Malace (USC postdoc) and Michael Paolone (USC grad. student)

Nucleon in the Nuclear Medium

- **Conventional Nuclear Physics:**
 - ▶ Nuclei are effectively and well described as point-like protons and neutrons (+ form factor) and interaction through effective forces (meson exchange)
 - ▶ Medium effects arise through **non-nucleonic degrees of freedom**
- **Nucleon Medium Modifications:**
 - ▶ Nucleons and mesons are not the fundamental entities in QCD
 - ▶ In the chiral limit, phase transition to quark-gluon plasma
 - ▶ Medium effects arise through **changes of fundamental properties of the nucleon**

The EMC Effect

- Depletion of the nuclear structure function $F_2^A(x)$ in the valence-quark regime $0.3 \leq x \leq 0.8$
- J. Smith and G. Miller: chiral quark-soliton model of the nucleon
Conventional nuclear physics does not explain EMC effect



J.R. Smith and
G.A. Miller, Phys.
Rev. Lett. **91**,
212301 (2003)

- → Nucleon structure is modified in the nuclear medium
- Note: prelim. E03-103 ^4He data consistent with SLAC A=12 param.

Limits for Medium Modifications

- Best constraints from **y-scaling**
 - ▶ $Q^2 > 1 \text{ (GeV/c)}^2$, $\Delta G_M < 3\%$ [1]
- **Coulomb Sum Rule**, L-Response
 - ▶ No quenching in the data observed [2]
 - ▶ Quenching of S_L is experimentally established [3]
 - ▶ Good agreement between theory and experiment for ${}^4\text{He}$ when using free-nucleon form factors [4]
 - ▶ $Q^2 \leq 0.5 \text{ (GeV/c)}^2$: $\Delta G_E < 15\%$ or even $< 5\%$

[1] I. Sick, Phys. Lett. B **157**, 13 (1985)


[2] J. Jourdan, Nucl. Phys. A **603**, 117 (1996)

[3] J. Morgenstern, Z.-E. Meziani, Phys. Lett. B **515**, 269 (2001)

[4] J. Carlson, J. Jourdan, R. Schiavilla, and I. Sick, Phys. Lett. B **553**, 191 (2003)


Quark Meson Coupling Model (QMC)

- **Structure of the nucleon** described by valence quarks in a bag (Cloudy-bag model).



intermediate baryon restricted to N or Δ

- **Nuclear system** described using effective scalar (σ) and vector (ω) meson fields.





- Scalar and vector fields of nuclear matter couple directly to confined quarks.

→ Modification of **internal structure** of bound nucleon

D.H. Lu, A.W. Thomas, K. Tsushima, A.G. Williams, K. Saito, Phys. Lett. B **417**, 217 (1998)
D.H. Lu *et al.*, Phys. Rev. C **60**, 068201 (1999)


Bound Proton EM Form Factors



- Electromagnetic rms radii and magnetic moment of the bound proton are increased
- **Charge form factor** much more sensitive to the nuclear medium than the **magnetic** ones.

D.H. Lu *et al.*, Phys. Rev. C **60**, 068201 (1999)



Chiral Quark Soliton Model (CQSM)



- **Chiral-soliton model** provides the quark and antiquark substructure of the proton, embedded in nuclear matter.
- Medium modifications:
 - ▶ significant for the ratio G_E/G_M
 - ▶ no strong enhancement of the magnetic moment

CQSM: J.R. Smith and G.A. Miller, Phys. Rev. C **70**, 065205 (2004)

Extended Skyrme Model



- Model of the nucleon based on **Skyrme Lagrangian**
- Results comparable to QMC, but differ in details
- $(G_E/G_M)_{\text{medium}}/(G_E/G_M)_{\text{free}} \approx 1$ for R = 1 fm

U. Yakhshiev, U. Meißner, A. Wirzba, Eur. Phys. J. A **16**, 569 (2003)

Other Models

- **Nambu–Jona-Lasinio model**

T. Horikawa, W. Bentz, Nucl. Phys. A **762**, 102 (2005)

- ▶ Nucleon as **quark-diquark** bound state + **nuclear matter** in the mean field approximation.
- ▶ Medium modifications: increase of the electric size in the medium
- ▶ **Medium modifications decrease with increasing Q^2** for both, spin and orbital form factors.

- **S. Liuti**

S. Liuti, hep-ph/0608251, hep-ph/0601125

- ▶ Connection between the modifications induced by the nuclear medium of the nucleon form factors and of the deep inelastic structure functions, obtained using the concept of **generalized parton distributions**.


Polarization-Transfer Technique

- **Free** electron-nucleon scattering

$$\frac{G_E}{G_M} = -\frac{P'_x}{P'_z} \cdot \frac{(E_i + E_f)}{2m} \tan\left(\frac{\theta_e}{2}\right)$$

- **Bound** nucleons → evaluation within model
Reaction-mechanism effects in $A(\vec{e}, e'\vec{p})B$
predicted to be small and minimal for
 - ▶ Quasielastic scattering
 - ▶ Low missing momentum
 - ▶ Symmetry about $\mathbf{p}_m = 0$

Proton Elastic Form-Factor Ratio




- Systematic decrease of G_E / G_M indicating difference in spatial distribution of charge and magnetization currents in the proton.
- Discrepancy can possibly be resolved by the inclusion of two photon effects in Rosenbluth analysis.

I.A. Qattan, Phys. Rev. Lett. **94**, 142301 (2005); Hall A E01-001

E93-049 and E03-104 at Jefferson Lab Hall A

${}^4\text{He}(e, e' \vec{p}){}^3\text{H}$ in quasielastic kinematics $Q^2 = 0.5 - 2.6 \text{ (GeV/c)}^2$



- Polarization-transfer ratio P'_x/P'_z : sensitive to G_E/G_M
- Induced polarization P_y : sensitive to final-state interactions



S. Dieterich, *et al.*, Phys. Lett. **B500**,47(2001); S. Strauch, *et al.*, Phys. Rev. Lett. **91**, 052301(2003); JLab E03-104, R.Ent, R. Ransome, S. Strauch, P. Ulmer (spokespersons)

Polarization Measurement


Focal-Plane Polarimeter



Observed angular distribution

$$\begin{aligned}
 I(\vartheta, \varphi) &= I_0(\vartheta) (1 + \epsilon_y \cos \varphi + \epsilon_x \sin \varphi) \\
 &= I_0(\vartheta) [1 + A_C (P_y \cos \varphi - P_x \sin \varphi)]
 \end{aligned}$$


Free Proton Form-Factor Ratio G_E/G_M



- Preliminary results from E03-104 with small statistical uncertainties $\delta(P'x/P'z) \approx 0.7\%$
- Full analysis of E03-104 will have reduced systematic uncertainties

${}^4\text{He}(\vec{e}, e' \vec{p})$ - Polarization-Transfer Ratio

$$R = P'_x/P'_z({}^4\text{He}) / P'_x/P'_z({}^1\text{H})$$




- RDWIA and RMSGA models can not describe the data.
- New data will set tight constraints

RDWIA: J.M. Udias *et al.*, Phys. Rev. Lett. **83**, 5451 (1999);


RMSGGA: P. Lava *et al.*, Phys. Rev. C **71**, 014605 (2005), D. Debruyne *et al.*, Phys. Rev. C **62**, 024611 (2000)

${}^4\text{He}(\vec{e}, e' \vec{p})$ - Polarization-Transfer Ratio




- $R^{\text{RDWIA}} \approx 0.97 \times R^{\text{RPWIA}}$
- Small sensitivity to
 - ▶ bound-state wave function
 - ▶ current operator
 - ▶ optical potential
- Enhancement of lower components (spinor distortions) in RDWIA

Polarization Transfer in ${}^4\text{He}(\vec{e}, e' \vec{p})$



- Previous data effectively described by **proton medium modified form factors**
- Preliminary data from E03-104 possibly hint at an unexpected trend in Q^2


Charge-Exchange FSI



- R suppressed by about 4% from MEC
- Spin-dependent charge exchange FSI suppresses R by about 6% and provides for alternative explanation
- CH-EX term not well constrained \Rightarrow need P_y from E03-104

R. Schiavilla *et al.*, Phys. Rev. Lett. **94**, 072303 (2005)


Induced Polarization in ${}^4\text{He}(e, e' \vec{p})$



- P_y is a measure of **final-state interactions**
- Observed final-state interactions are small and increase with missing momentum
- RDWIA results consistent with data
- RDWIA can be used to correct data for HRS acceptance (30% - 40% effect)

E03-104 induced polarization still very preliminary.
Uncertainties are statistical only; systematic uncertainties < 0.02

Induced Polarization in ${}^4\text{He}(e, e' \vec{p})$




- Observed final-state interaction small and with **very weak Q^2 dependence**
- RDWIA results consistent with data
- Spin-dependent charge exchange terms not constrained by N-N scattering and possibly overestimated
- E03-104 took specific data that will set tight constraints on FSI


Inner uncertainties are statistical only;
full analysis of E03-104 will have reduced systematic uncertainties

New Proposal to Measure ${}^4\text{He}(e,e'p){}^3\text{H}$ in Hall C

- Proton spectrometer: HMS
Electron spectrometer: SOS
(momentum bite: 40%)
- HMS is being equipped with FPP for G_E^p -III
- **Missing mass resolution sufficient** to identify ${}^3\text{H}$ in the final state if SOS is used as electron spectrometer.




Q^2 Distribution of R




- Anticipated data in 27 days of beam time on ^4He

Missing-Momentum Distribution



- Improved constraints on models through missing-momentum distribution
- “Would be nice to study modification of the nucleon form factors as a function of the nucleon momentum.” [\[Mark Strikman\]](#)

Missing-Momentum Distribution



- Proposed Hall C data: induced polarization

Summary

- **Proton in the nuclear medium**
 - ▶ Models predict change of the internal structure of a bound nucleon
 - ▶ Corrections due to in-medium form factors could be significant
- **Polarization transfer in $^4\text{He}(e,e'p)$**
 - ▶ Significant deviation from RDWIA results; data **effectively described by proton medium modifications**
 - ▶ Alternative interpretation in terms of strong **charge-exchange FSI**
 - ▶ Induced polarization crucial to clarify role of FSI
 - ▶ New results from E03-104 will provide needed constraints
 - ▶ Experiment in Hall C could measure missing-momentum distributions and extend the data set to larger Q^2