



The Continuous Electron Beam Accelerator Facility
Theory Group Preprint Series

Additional copies are available from the authors.

The Southeastern Universities Research Association (SURA) operates the Continuous Electron Beam Accelerator Facility for the United States Department of Energy under contract DE-AC05-84ER40150

CEBAF-TH-96-09
DOE/ER/40561-258-INT96-19-02
April 22, 1996

Interaction of small size wave packet with hadron target

L. FRANKFURT*

*School of Physics and Astronomy
Raymond and Beverly Sackler Faculty of Exact Sciences
Tel Aviv University 69978
Tel Aviv, Israel*

A. RADYUSHKIN†

*Physics Department, Old Dominion University, Norfolk, VA 23529, USA
and
Continuous Electron Beam Accelerator Facility,
Newport News, VA 23606, USA*

M. STRIKMAN‡

*Department of Physics, Pennsylvania State University
University Park, PA 16802, USA*

Abstract

We calculate in QCD the cross section for the scattering of an energetic small-size wave packet off a hadron target. We use our results to study the small- σ behaviour of $P_\pi(\sigma)$, the distribution over cross section for the pion, in the leading α_s -order.

DISCLAIMER

This report was prepared as an account of work sponsored by the United States government. Neither the United States nor the United States Department of Energy, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, mark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or any agency thereof.

*On leave of absence from the St.Petersburg Nuclear Physics Institute, Russia

†Also Laboratory of Theoretical Physics, JINR, Dubna, Russian Federation

‡Also St.Petersburg Nuclear Physics Institute, Russian Federation

1. INTRODUCTION

Recently phenomena involving interactions of hadrons in small-size configurations have been intensively discussed both in relation with the phenomenon of color transparency and vector meson electroproduction observed at HERA energies. There is also a deep relation between presence of the weakly interacting small size configurations in hadrons and phenomenon of cross section fluctuations in the interactions of the hadrons which manifests itself in the inelastic coherent diffraction processes: $h + N(A) \rightarrow X + N(A)$, see Ref. [1]. In this paper, we focus on the systematic derivation of the formulae for the interaction of the color singlet $q\bar{q}$ pair having a small transverse size with a hadron target. Then, we use these formulae to calculate the probability of the distribution for the interaction of a photon and a pion with a target for small interaction cross sections. Although some of equations deduced in the paper existed before no derivations with analysis of their accuracy has been presented.

The paper is organized as follows. In Section 2, we consider the virtual forward Compton amplitude in the small- x region where it is dominated by the photon-gluon scattering subprocess. We outline there a derivation of the basic formula expressing the total cross section σ_{γ^*T} as a convolution of the gluon distribution amplitude $G_T(x, Q^2)$ and the γg scattering cross section. In Section 3, we write down the γg cross section in terms of the $\bar{q}q$ light-cone wave functions of the virtual photons. In the next section, we calculate the cross section distribution $P_{\gamma^*}(\sigma)$ for the virtual photon. In Section 5, we discuss the quark-hadron duality interplay between the perturbative free-quark results and contributions due to low-lying resonances. Finally, in Section 6, we calculate the cross section distribution for the pion $P_\pi(\sigma)$ in the small cross section limit where it is governed by $\bar{q}q$ configurations having small spatial size. Basing on QCD evolution equation we evaluate also the functional dependence of $P_\pi(\sigma) \rightarrow 0$ on σ and on the incident energy.

2. HARD γ^*T TOTAL CROSS SECTION AND THE INTERACTION OF SMALL SIZE CONFIGURATIONS.

Let us consider a particular contribution into the γ^*T cross section corresponding to a transformation of the virtual photon when γ^* converts into a $Q\bar{Q}$ pair with quarks having a large relative transverse momentum. Usually, this contribution is written as a convolution of the infinite momentum frame wave function of the target with the pQCD calculable coefficient function describing

the short-distance propagation of the particles between two virtual photon vertices. Our aim is to express the relevant coefficient function in terms of the light-cone wave functions of the virtual photons as viewed from the reference frame where the target is at rest. The contribution we are interested in is given by the sum of diagrams shown in Fig.1.

The lower blob corresponds to the gluon distribution in the target. It is convenient to parameterize the gluon momentum k in terms of the Sudakov variables

$$k = -\alpha q' + \beta p' + k_t \quad d^4k = \frac{s}{2} d\alpha d\beta d^2k_t. \quad (2.1)$$

Here q' and p' are light-like momenta related to p, q by

$$q = q' + \frac{q^2}{2(p'q')}p' ; \quad p = p' + \frac{p^2}{2(p'q')}q'; \quad (2.2)$$

$$2(pq) = 2(p'q') + \frac{q^2p^2}{2(p'q')}. \quad (2.3)$$

For our goals, the most interesting region is that of small values of the Bjorken parameter: $x = -q^2/2(pq) \rightarrow 0$, where we may safely approximate $2(p'q') = s$.

In the d^4k integral, the region $k_t^2 \sim Q^2$ corresponds to the next-order α_s correction, so we will take into account only the contribution of the region $k_t^2 \ll Q^2$. This corresponds to the leading $\alpha_s \log Q^2/\Lambda^2$ approximation in which the $\mathcal{O}(\alpha_s)$ corrections are neglected. In this kinematical region, the contribution of the diagrams shown in Fig. 1 can be considerably simplified. The essential region of integration is

$$|k^2| = |\alpha\beta s + k_t^2| \ll Q^2. \quad (2.4)$$

Note that $\beta s \sim Q^2 \propto (mass)^2$ of the $q\bar{q}$ state produced by γ^* . Hence,

$$\alpha \ll 1 \quad (2.5)$$

is the essential region of integration over α . As a result, it is sufficient to take into account only the longitudinal polarization in the propagator of the exchanged gluon [2]

$$\frac{d_{\mu\lambda}(k)}{k^2} \simeq \frac{p'_\mu q'_\lambda}{(p'q')} \frac{1}{k^2}. \quad (2.6)$$

Indeed, let us estimate the contribution due to exchange of a transversely polarized gluon:

$$\delta\sigma \sim \frac{1}{(2\pi)^4} \int \frac{T_{\mu\perp\mu\perp}^{\gamma^*}}{s} d(\beta s) \int \frac{T_{\bar{\mu}\perp\bar{\lambda}\perp}^T}{s} d(\alpha s) \frac{d^2 k_t}{(k^2)^2}. \quad (2.7)$$

Here, $T_{\mu\perp\lambda\perp}^{(\gamma^*)}$ is the imaginary part of the amplitude of the γ^* scattering off a gluon given by the lowest-order Feynman diagrams and $T_{\bar{\mu}\perp\bar{\lambda}\perp}^T$ is that for the gluon scattering off a target T .

Using the fact that, at high energies, the Feynman amplitude of processes due to exchange by two elementary fermions tends to constant [3] we obtain:

$$\int T_{\lambda\perp\lambda\perp}^{\gamma^*} d(\beta s) \propto \frac{(\beta s)^2}{Q^2} \simeq (Q^2).$$

In this estimate we use also scaling over Q^2 in the box diagram. The amplitude due to the exchange by two vector particles increases like s , and we have

$$\int T_{\mu\perp\lambda\perp}^{(T)} d\alpha s \propto (\alpha s)^{2+n}.$$

Here $n > 0$, since, according to the QCD evolution equations, the deep inelastic amplitudes increase with energy in region of applicability of the perturbative QCD.

As a result of this power counting estimate we obtain:

$$\begin{aligned} \delta\sigma &\sim \int \frac{(\beta s)^2}{s} \frac{(\alpha s)^{2+n}}{s} \frac{d^2 k_t}{(\alpha\beta s + k_t^2)^2} \theta(k_t^2 < k_{i0}^2) \theta(\alpha\beta s \leq k_{i0}^2) \\ &\approx \int \frac{k_{T0}^4 d^2 k_t}{k_i^4} \theta(k_t^2 < k_{i0}^2) (\alpha s)^n \propto (\alpha s)^n. \end{aligned} \quad (2.8)$$

Here, we substituted $\alpha\beta s \sim k_i^2$. The divergence at small k_t disappears when matrix element is calculated more accurately. Thus, due to the presence of the factor α in eq.(2.8), in the leading $\alpha_s \log Q^2/\lambda^2$ approximation, the contribution due to the exchange of a transversely polarized gluon is negligible compared to the contribution of the longitudinal polarization specified by eq.(2.6). We use here the observation that in QCD the power n characterizing the energy dependence of the amplitude is the same for scattering of transversely and longitudinally polarized gluons.

Using the gluon propagator in the form given by eq.(2.6), we get the following expression for the total contribution of the diagrams shown in Fig. 1:

$$\text{Im } M = \int \frac{d^4 k}{(2\pi)^4 i} \frac{1}{(k^2)^2} 2 \cdot \text{Im } T_{\mu\lambda}^{ab(P)} \text{Im } T_{\mu\lambda}^{ab(T)} \quad (2.9)$$

Here $T_{\mu\lambda}^{ab(P)} = T_{\mu\lambda}^{ab}(\gamma^* g \rightarrow Q\bar{Q})$ is the sum of the box diagrams describing the $\gamma^* g$ scattering and $T_{\mu\lambda}^{ab(T)}$ is the amplitude of the gluon scattering off the target T . We use here the fact that

$$T_{\mu\lambda}^{ab(P)} k_\mu = T_{\mu\lambda}^{ab(P)} k_\lambda = 0, \quad (2.10)$$

since the box diagram contains no gluons and, therefore, the Ward identities in this approximation are the same as in an Abelian gauge theory (a, b are the color indices).

Using the dominance of the longitudinal gluon polarization (2.6) and incorporating eq.(2.1) we can simplify eq.(2.9):

$$\frac{\text{Im } M}{s} = \int \frac{sd\alpha d\beta d^2 k_t}{2(2\pi)^4 (k^2)^2} \frac{2\text{Im } T_{\mu\lambda}^{ab(P)} p_\mu p_\lambda}{4(pq)^2} \cdot \frac{4\text{Im } T_{\bar{\mu}\bar{\lambda}}^{ab(T)} q_\mu q_\lambda}{s}. \quad (2.11)$$

From eq.(2.1) and (2.10) it follows that

$$\frac{\text{Im } T_{\mu\lambda}^{ab(P)} p_\mu p_\lambda}{4(pq)^2} = \frac{\text{Im } T_{\mu\perp\lambda\perp}^{ab(P)} k_t^\mu k_t^\lambda}{(\beta s)^2} \quad (2.12)$$

and

$$\frac{\text{Im } M}{s} = \int d\beta s \frac{\frac{1}{2} \sum T_{\mu\perp\mu\perp}^{ab(P)}}{(\beta s)^2} \int \frac{d\alpha s d^2 k_t}{(2\pi)^4 (k^2)^2} \frac{4k_t^2 \text{Im } T_{\bar{\mu}\bar{\lambda}}^{ab(T)} q_\mu q_\lambda}{s^2} \quad (2.13)$$

It is useful to define the cross section of γ^* scattering off a gluon g averaged over the gluon color:

$$\delta_{ab} \cdot s' \cdot \sigma(\gamma^* g \rightarrow q\bar{q}) = \frac{1}{2} \sum_{\mu\perp=1,2} \text{Im } T_{\mu\perp\mu\perp}^{ab(P)}, \quad (2.14)$$

where s' is the invariant mass of the produced $q\bar{q}$ system: $s' = (k+q)^2 \simeq \beta s - Q^2$. Thus,

$$\sigma_{\gamma^* T} = \frac{\text{Im } M}{s} = \int \frac{d\beta}{\beta} \sigma(\gamma^* g \rightarrow q\bar{q}) \int \frac{sd\alpha d^2 k_t}{(2\pi)^4 (k^2)^2} k_t^2 \sum_a \frac{4\text{Im } T_{\bar{\mu}\bar{\lambda}}^{a(T)} q_\mu q_\lambda}{s^2}. \quad (2.15)$$

In the leading $\alpha_s \log Q^2$ approximation, we can substitute k^2 by k_t^2 . Comparing our result with the QCD-improved parton model expression for the production of heavy quarks (see *e.g.*, [4]), we observe that

$$\int \frac{sd\alpha d^2k_t}{(2\pi)^4 k_t^2} \sum_a 4\text{Im} T_{\mu\lambda}^{aa(T)} q_\mu q_\lambda = x G_T(x, Q^2), \quad (2.16)$$

where G_T is the gluon distribution in a target T . This gives

$$\sigma_{\gamma^* T} = \int \sigma_{\gamma^* g} \frac{d\beta}{\beta} [x G_T(\beta, Q^2)]. \quad (2.17)$$

As the first argument of $\beta G(\beta, Q^2)$, we use $\beta = \frac{Q^2 + M^2}{s} \approx \frac{2Q^2}{s} \approx 2x$ Here M is the mass of produced $q\bar{q}$ pair. As the evolution scale for the gluon distribution function, we take Q^2 . Of course, the higher-order α_s corrections may change Q^2 by some numerical factor. This scale-fixing ambiguity is a usual feature of the leading $\alpha_s \log Q^2$ -approximation.

3. LIGHT-CONE WAVE FUNCTIONS AND $\sigma_{\gamma^* G}$

Now let us express $\sigma_{\gamma^* g}$ in terms of the light-cone wave functions of the virtual photon. To this end, we write down the four-momenta $r_1(r_2)$ of quark (antiquark) in the box in terms of the light-cone variables $r_1 = \{r_1^+, r_1^-, r_t\}$ with $r_1^+ = \eta q^+$ and take the integral over r_1^- by residue. Introducing the lowest-order perturbative $q\bar{q}$ light-cone wave functions of the virtual photon [5]

$$\psi_\mu = \frac{\bar{U}(r_1)\gamma_\mu U(-r_2)}{\frac{m^2 + r_t^2}{\eta(1-\eta)} + Q^2} \frac{1}{\sqrt{\eta(1-\eta)}}, \quad (3.18)$$

we obtain the following expression for the sum of the box diagrams:

$$\int d\alpha s \frac{\text{Im} T_{\mu\lambda}^{ab(p)}}{s^2} p_\mu p_\lambda = e^2 g_s^2 \int \frac{d\eta d^2 r_t}{2(2\pi)^3} \pi \psi_\mu(\eta, r_t) \{2\psi_\mu(\eta, r_t) - \psi_\mu(\eta, r_t + k_t) - \psi_\mu(\eta, r_t - k_t)\} F_a F_b, \quad (3.19)$$

where g_s^2 is the QCD coupling constant and $F_a = \frac{\lambda_a}{2}$, λ_a being the Gell-Mann matrices of the $SU(3)_c$ group in the fundamental representation.

It is convenient to rewrite this formula in the impact parameter space:

$$\psi_\mu(x, r_t) = \int \psi_\mu(x, b) e^{i r_t \cdot b} d^2 b. \quad (3.20)$$

Then

$$\int d\alpha s \frac{\text{Im} T_{\mu\lambda}^{PP} p^\mu p^\lambda}{(2pq)^2} = \int \psi_\mu^2(x, b) \frac{dx d^2 b}{4\pi} g_s^2 \{ \pi [2 - e^{i k_t \cdot b} - e^{-i k_t \cdot b}] \text{Tr}(F_a F_b) \}. \quad (3.21)$$

Within the leading $\alpha_s \log Q^2$ approximation, to obtain eq.(2.12), it is necessary to decompose exponent into a power series over $(k_t b)$ and to keep terms up to the second order in k_t^2 . Combining eqs. (3.20), (2.11), and (2.16), we obtain

$$\sigma_{\gamma^* T} = e^2 \int \psi_\mu^2(\eta, b) \frac{dz d^2 b}{4\pi} N_c \left\{ \frac{1}{N_c} g_s^2 \pi \frac{(k_t b)^2}{k_t^2} \text{Tr} \frac{F^2}{8} \right\} G_T(x, \lambda/b^2). \quad (3.22)$$

Here, factor λ can be estimated from analysis of $\sigma_L(\gamma^* N)$ cross section. Since the gluon density increases when x decreases, λ slowly increases with decrease of x [6]. For $x \sim 10^{-3}$, $\lambda \approx 9$.

It is instructive to represent $\sigma_{\gamma^* T}$ in the form

$$\sigma_{\gamma^* T} = e^2 \int \psi_\mu^2(\eta, b) \frac{d\eta d^2 b}{4\pi} N_c \cdot \sigma_i^{q\bar{q}}(b^2). \quad (3.23)$$

Here, $\sigma_i^{q\bar{q}}$ is the cross section for the interaction of a colorless small transverse size $q\bar{q}$ -pair with the target T :

$$\sigma_i^{q\bar{q}} = g_s^2 \pi \frac{b^2}{2} \frac{1}{N_c} \text{Tr} \left(\frac{F^2}{8} \right) x G_T(x, \lambda/b^2). \quad (3.24)$$

This expression was obtained originally in [7,8]. As usual, N_c is the number of colors, and the Casimir operator of the $SU(3)$ group in the fundamental representation can be easily calculated:

$$\frac{1}{8} \frac{1}{N_c} \text{Tr} F^2 = \frac{1}{3} \text{Tr} F_3^2 = \frac{1}{6}. \quad (3.25)$$

Combining all the numbers together, we finally obtain:

$$\lambda \sigma_i^{q\bar{q}} = \frac{\pi^2}{3} b^2 [x G_T(x, \lambda/b^2)] \alpha_s(\lambda/b^2). \quad (3.26)$$

Here $b = (b_q - b_{\bar{q}})$. This formula describes the essence of the color transparency (CT) phenomenon (*cf.* discussion in [9]): $q\bar{q}$ configuration of a small spatial size has a small interaction cross section. However, for sufficiently small x , the interaction becomes strong due to the formation of the soft gluon field. In this respect, eq.(3.22) predicts the interaction of a small size configuration which is

qualitatively different from that of the models of F. Low [10] and J. Gunion and D.Soper [11]. The fact that $\sigma_T^{q\bar{q}}$ is proportional to the gluon distribution in eq.(3.24) increasing in the small- x region, has important experimental consequences, *e.g.* it makes it possible to observe the small-size quark configurations at HERA in the electroproduction of vector mesons at small x . In fact, eq.(3.22) can be inferred from a formula derived in ref. [12] within a model approximation to QCD. Using some simple tricks, one can also obtain eq.(3.22) from a formula obtained in [13] within the leading $\alpha_s \ln x$ approximation of QCD combined with some bold assumptions concerning the parton model structure.

Using (3.19), we can calculate distribution over cross section for the fast photon or pion projectible for small σ (*cf.* [7]).

4. DISTRIBUTION OF P_{γ^*} FOR THE PHOTON PROJECTILE.

In the previous section we have derived eq.(3.22) which expresses the σ_{γ^*T} cross section in terms of the light-cone wave functions of the virtual photon γ^* . This formula gives us the possibility to calculate another useful quantity - distribution over cross section. It has been understood long ago [14–16] that many features of the interaction of a fast projectile can be described in terms of distribution over cross section. An important advantage of such a quantity is that it accurately takes into account diffractive processes. Properties of $P(\sigma)$ have been discussed in detail in [17]. So we restrict our consideration by a calculation of $P_{\gamma^*}(\sigma)$ at $\sigma \rightarrow 0$.

By definition, the experimentally observable total cross section in terms of $P(\sigma)$ is given by

$$\sigma_{\gamma^*N} = \int P_{\gamma^*}(\sigma) \sigma d\sigma. \quad (4.1)$$

Incorporating here eq.(3.24), we can write

$$e\sigma_{\gamma^*N} = e^2 \int \psi_{\gamma^*}^2(\eta, b) \frac{d\nu}{4\pi} N_c \sigma \frac{\pi db^2}{d\sigma}. \quad (4.2)$$

Since eq.(3.24) is applicable for small σ only, comparing eqs.(4.1) and (4.2), we obtain

$$P_{\gamma^*}(\sigma \rightarrow 0) = \int e^2 \psi_{\gamma^*}^2(\nu, b) \frac{d\nu}{4} N_c \frac{\pi db^2}{d\sigma}, \quad (4.3)$$

where $\psi_{\gamma^*}(\eta, b)$ is defined by eqs.(3.18) and (3.20). The functional dependence of b on σ in eq. (2.3) should be calculated from eq.(3.24).

Eq. (4.3) predicts a rather complicated dependence of $P(\sigma)$ on $\ln \sigma$ at small σ . However, this dependence can be easily calculated using QCD evolution equations. The distinctive feature of eq.(4.3) is that

$$P_{\gamma^*}(\sigma \rightarrow 0) |_{\sigma \rightarrow 0} \sim \frac{1}{\sigma} \quad \text{up to } \log(\sigma/\sigma_0) \text{ terms.} \quad (4.4)$$

5. TRANSITION TO MESONS

The virtual photon wave function $\psi_\mu(\eta, r_t)$ (1.19) can be written through a dispersion integral

$$\psi_\mu(\eta, r_t) = \frac{1}{\pi} \int_0^\infty \psi_\mu^{\bar{q}q}(\kappa; \eta, r_t) \frac{d\kappa^2}{\kappa^2 + Q^2} \quad (5.1)$$

where

$$\psi_\mu^{\bar{q}q}(\kappa; \eta, r_t) = \frac{\bar{U}(\eta r_+) \gamma_\mu U((1-\eta)r_+)}{\sqrt{\eta(1-\eta)}} \delta\left(\kappa^2 - \frac{m_q^2 + r_t^2}{\eta(1-\eta)}\right) \quad (5.2)$$

is the wave function of a non-interacting $\bar{q}q$ -pair with invariant mass κ . Due to interaction between the quarks, the spectral density of the dispersion representation (5.1) is modified and, instead of the free-quark approximation $\psi_\mu^{\bar{q}q}(\eta, \kappa)$, one should use a sum over resonances, the ρ -meson being the dominant feature in the low- κ region:

$$\psi_\mu^{\bar{q}q}(\eta, \kappa) \rightarrow g_\rho \psi_\mu^\rho(\eta) \delta(\kappa^2 - m_\rho^2) + \psi_\mu^{\text{higher states}}(\eta, \kappa) \quad (5.3)$$

where g_ρ is the magnitude of the ρ -state projection onto the electromagnetic current. At large κ , the resonances are wide, and their sum rapidly approaches the free-quark value, *i.e.*, one has a perfect quark-hadron duality. For sufficiently large Q^2 , the dispersion integral is dominated by higher states, and the free-quark approximation is completely justified. Decreasing Q^2 , one would observe mismatch between the free-quark calculation and the dispersion integral over the resonances. Such a situation is well known from QCD sum rules: the difference between the resonance and free-quark spectra is described by power corrections

$(1/Q^2)^N$. The usual procedure is to approximate the higher states by the free-quark contribution (“first resonance plus continuum” model)

$$\psi_\mu^{\text{higher states}}(\eta, \kappa) = \theta(\kappa^2 > s_0^\rho) \psi_\mu^{\bar{q}q}(\eta, \kappa)$$

where s_0^ρ is the effective threshold for higher resonances in the ρ -channel and then fix its value by the requirement of the best agreement between the two sides of the resulting sum rule

$$\frac{1}{\pi} \int_0^{s_0^\rho} (\pi g_\rho \psi_\mu^\rho(\eta) \delta(\kappa^2 - m_\rho^2) - \psi_\mu^{\bar{q}q}(\eta, \kappa)) \frac{d\kappa^2}{\kappa^2 + Q^2} = \sum_{N=2} \frac{A_N}{(Q^2)^N}. \quad (5.4)$$

After fixing s_0^ρ from the magnitude of the power corrections $A_N/(Q^2)^N$, one can take the limit $Q^2 \rightarrow \infty$ to get the local duality relation

$$\pi g_\rho \psi_\mu^\rho(\eta) = \int_0^{s_0^\rho} \psi_\mu^{\bar{q}q}(\eta, \kappa) d\kappa^2. \quad (5.5)$$

In other words, the ρ -meson wave function in such an approach is dual to the free-quark wave functions integrated over the duality interval $0 \leq \kappa^2 \leq s_0^\rho$.

For the forward virtual Compton amplitude, the dispersion representation can be applied both for the initial and “final” virtual photon. Taking only the ρ -meson contribution in both dispersion integrals one would get the amplitude for the $\rho T \rightarrow \rho T$ scattering. This idea can be also used to study the pion diffractive scattering.

6. CALCULATION OF $P_\pi(\sigma \rightarrow 0)$.

To analyze the pion scattering, we substitute the electromagnetic current by the axial current in the original amplitude, *i.e.*, simply add γ_5 in the current vertices. For massless quarks, the final result has the same structure as that for the vector current. Of course, the $\bar{q}q$ -pair wave function would have an extra γ_5 . The vertex function for the transition of the axial current into quarks with four momenta r_1 and r_2 is

$$\frac{\bar{U}(r_1) \gamma_\mu \gamma_5 U(-r_2)}{\sqrt{x(1-x)}} \simeq 2(r_1 + r_2)_\mu. \quad (6.1)$$

The projection of a single-pion state onto the axial current is specified by the $\pi \rightarrow \mu\nu$ decay constant f_π :

$$\langle 0 | J_\mu^A | \pi, P \rangle = \sqrt{2} f_\pi P_\mu. \quad (6.2)$$

Hence, we should take the amplitude $\sim P_\mu P_\nu$ corresponding to the longitudinal polarization of the axial current. Again, the transition from the virtual amplitude for the currents to that involving the pion can be understood in terms of the dispersion representation and quark-hadron duality. In other words, below the effective higher state threshold s_0^π , one should substitute the free-quark contribution by that due to the pion pole:

$$\psi_{5\mu}^{\bar{q}q}(\eta, \kappa) \rightarrow q_\mu (f_\pi \psi_\pi(\eta) \delta(\kappa^2 - m_\pi^2) + \theta(\kappa^2 > s_0^\pi) \psi_\mu^{\bar{q}q}(\eta, \kappa)). \quad (6.3)$$

The local duality prescription gives a correctly normalized wave function provided that $s_0^\pi = 16\pi^2 f_\pi^2 \approx 0.67 \text{ GeV}^2$. Of course, one can use a pion wave function different from that given by the local duality. However, the duality considerations justify the use of the effective two-body wave function (see [18]).

The actual calculation consists of the same steps as those leading to eq.(3.25). For a small-size configuration, we get the following contribution $\delta\sigma_{\pi N}$ into the scattering cross section $\sigma_{\pi\mu}^{\text{tot}}$:

$$\delta\sigma_{\pi N} = \int |\psi_\pi(\eta, b)|^2 \frac{d\eta d^2b}{4\pi} N_c \sigma_N^{q\bar{q}}(b^2). \quad (6.4)$$

Effectively, the vertex $e\psi_{\gamma^*} \cdot \sqrt{N_c}$ is substituted by the pion wave function. Rewriting $\delta\sigma_{\pi N}$ as

$$\delta\sigma_{\pi N} = P_\pi(\sigma) d\sigma = \frac{db^2}{d\sigma} \int |\psi_\pi(\eta, b)|^2 \frac{d\eta}{4} \sigma d\sigma, \quad (6.5)$$

we obtain:

$$P_\pi(\sigma \rightarrow 0) = \frac{db^2}{d\sigma} \int |\psi_\pi(\eta, b \rightarrow 0)|^2 \frac{d\eta}{4}, \quad (6.6)$$

where $\sigma(b^2)$ is given by eq.(3.26).

Thus, $P_\pi(\sigma \rightarrow 0)$ is determined by the pion wave function at the origin of the impact parameter space, or, what is the same, by the integral of the momentum wave function $\psi_\pi(\eta, r_i)$ over all transverse momenta r_i . This integral formally gives the pion distribution amplitude

$$\varphi_\pi(\eta) = \frac{\sqrt{3}}{(2\pi)^3} \int \psi_\pi(\eta, r_i) d^2r_i.$$

However, in QCD (and in any theory with dimensionless coupling constant), this integral diverges. The standard procedure is to supplement the integral with some renormalization prescription characterized by a cut-off parameter μ , *i.e.*, $\varphi_\pi(\eta) \rightarrow \varphi_\pi(\eta, \mu)$. In fact, the Fourier transformation from the momentum to the impact parameter space

$$\psi_\pi(\eta, b) = \int \psi_\pi(\eta, r_t) \frac{d^2 r_t}{(2\pi)^2}$$

for small b can also be treated as a particular cut-off prescription with $1/b$ playing the role of the renormalization parameter μ . In the $b \rightarrow 0$ limit, one encounters the singular $\log(b^2)$ terms. It is exactly the logarithms which generate the evolution of the pion distribution amplitude. Summing the logarithms by the renormalization group methods gives, for small b :

$$\psi_\pi(\eta, b) = \eta(1-\eta) \sum_{n=0} a_n C_n^{3/2}(2\eta-1) \left(\frac{\log b_0^2 \Lambda^2}{\log b^2 \Lambda^2} \right)^{\gamma_n/2\beta_0} \quad (6.7)$$

where $\eta(1-\eta)C_n^{3/2}(2\eta-1)$ are the eigenfunctions of the evolution kernel ($C_n^{3/2}(2\eta-1)$ being the Gegenbauer polynomials), the anomalous dimensions γ_n are its eigenvalues and β_0 is the one-loop QCD β -function coefficient. The b_0 -parameter characterizes the onset of the perturbative evolution. The coefficients a_n are the Gegenbauer moments of the pion wave function at this scale. Note that the anomalous dimension of the axial current vanishes ($\gamma_0 = 0$) and all other γ_n 's are positive. Hence, after the renormalization group improvement, the limit $b \rightarrow 0$ is well-defined in this case and

$$\psi_\pi(\eta, b=0) = \sqrt{48\pi} f_\pi \eta(1-\eta), \quad (6.8)$$

where $f_\pi = 92\text{MeV}$. The absolute normalization of the pion wave function for $b=0$ is fixed by the matrix element of the axial current:

$$\int \psi_\pi(\eta, r_t) \frac{d\eta d^2 r_t}{8\pi^3} = \frac{f_\pi}{\sqrt{N_c}}, \quad (6.9)$$

or in the impact parameter space (see eq.(3.20))

$$\int \psi_\pi(\eta, b=0) \frac{d\eta}{2\pi} = \frac{f_\pi}{\sqrt{N_c}}. \quad (6.10)$$

In other words, for the pion, the singular $\log b$ terms sum into harmless $(1/\log b^2 \Lambda^2)^{\gamma_n/2\beta_0}$ factors vanishing in the $b \rightarrow 0$ limit. As a result, the η -dependence of the pion wave function $\psi_\pi(\eta, b)$ in the formal $b \rightarrow 0$ limit always

assumes its asymptotic form $\psi_\pi(\eta, b) \sim \eta(1-\eta)$, irrespectively of its shape at the scale b_0 . It is natural to expect that b_0 is related to the scale characterizing the magnitude of the nonperturbative momentum distribution in the pion. The momentum scale $\mu_0 = \sqrt{s_0^2} \approx 0.8\text{GeV}$ suggested by the local duality is rather large, and there may exist a transitional region of distances $b \sim b_0$ small compared to the pion size but not small enough to produce sizable perturbative evolution effects. In this case, one can try the η -dependences of $\psi_\pi(\eta, b)$ different from the asymptotic form. In fact, the integral

$$I \equiv \int |\psi_\pi(\eta, b)|^2 \frac{d\eta}{4} \quad (6.11)$$

is rather insensitive to the evolution effects. If we take the asymptotic wave function (6.8) then

$$P_\pi(\sigma \rightarrow 0) = \frac{2}{5} \pi^2 f_\pi^2 \frac{db^2}{d\sigma}. \quad (6.12)$$

Assuming that, at the scale $b = b_0$, the η -dependence of the pion wave function corresponds to the Chernyak-Zhitnitsky [19] ansatz

$$\psi_\pi^{CZ}(\eta, b = b_0) = 5\sqrt{48\pi} f_\pi \eta(1-\eta)(1-2\eta)^2, \quad (6.13)$$

we obtain

$$P_\pi(\sigma, b = b_0) = \frac{10}{21} \pi^2 f_\pi^2 \frac{db^2}{d\sigma}. \quad (6.14)$$

Thus, in this case the evolution would decrease the integral I by $\sim 20\%$ when b changes from b_0 to 0.

Taking the asymptotic result, we get:

$$P_\pi(\sigma \rightarrow 0) = \frac{6}{5} \frac{f_\pi^2}{\alpha_s x G_N(x, \lambda/b^2)}. \quad (6.15)$$

Distribution $P_{\pi N}(\sigma)$ was determined in Ref. [7] from the analysis of the soft diffractive processes for $E_\pi \approx 200\text{GeV}$, see solid curves in Fig.2. In the limit $\sigma \ll \langle \sigma \rangle$, we can compare this result with eq.(6.14). The applicability region of this equation is restricted by several conditions. First, x_{eff} should be small enough so that the average longitudinal distances in the scattering process $1/2m_N x$ were larger than the nucleon size, which corresponds to $x < 0.05$. Furthermore, the virtualities in the process should be large enough so that one can apply pQCD which corresponds to the requirement $Q_{eff}^2 \geq 1 - 2\text{GeV}^2$. In our analysis we

also neglect the b -dependence of the wave function of the $q\bar{q}$ component (this is a higher twist effect), which restricts consideration to $b \leq 0.5 fm$. In the numerical calculation, we use GRV parameterization [20] since it describes well the parton distributions down to $Q^2 \sim 1.5 GeV^2$. We present results both for the leading and next-to-leading order GRV parameterizations, see dashed curves in Fig.2. Difference between LO and NLO results illustrates range of uncertainties of the current analysis. One can see that the results of our calculations are in qualitative agreement with the phenomenological results of [7].

Another interesting feature of our results is a substantial energy dependence of $P(\sigma < \langle \sigma \rangle)$ on the incident energy due to a fast increase of $xG_N(x, Q^2)$ with the decrease of x , see Fig.3. This reflects the fact that the probability of point-like configurations in hadrons decreases with the increase of energy. Further diffractive data (preferably at higher energies) are necessary to get better information about $P_{\pi N}(\sigma)$.

Since the existence of configurations with small spatial size has been confirmed experimentally in the energy dependence and absolute value of cross section of electroproduction of vector mesons, we consider the above result as a reflection of soft matching between nonperturbative and pQCD regimes.

7. SUMMARY AND CONCLUSIONS

In this paper, we applied a pQCD approach to describe the basic features of the high-energy interactions of a small-size $\bar{q}q$ configurations with a hadron target. This interaction is proportional to the gluon distribution function $G_T(x, Q^2)$, of the target and, hence, the cross section is enhanced in the small- x region. The $\bar{q}q$ configuration can be described by the wave functions whose particular form is determined by the projection of the initial particle (γ^* , ρ or π) onto the $\bar{q}q$ component. For small σ , we calculated the cross section distribution $P_{\pi}(\sigma)$ for the pion and demonstrated that it is rather insensitive to the specific form of the pion distribution amplitude.

8. ACKNOWLEDGMENTS

This work was partially supported by the BSF Grant No. 9200126 and by the US Department of Energy under contracts DE-FG02-93ER40771 and DE-AC05-84ER40150. Two of us (AR and MS) thank the DOE's Institute for Nuclear Theory at the University of Washington for its hospitality and support during the workshop "Quark and Gluon Structure of Nucleons and Nuclei".

REFERENCES

- [1] L. L. Frankfurt, G. A. Miller and M. Strikman, Ann. Rev. of Nucl. and Particle Phys. 44 (1994) 501.
- [2] V.N.Gribov, Lectures on the theory of complex momenta. KhFTI-70/48, Kharkov (1970).
- [3] Ya.I.Azimov Phys.Lett., 3,195(1963).
- [4] M.A. Shifman, A.I. Vainshtein, V.I. Zakharov Nucl.Phys.B136,157(1978); M. Gluck, E. Reya, Phys.Lett.83B, 98,(1979).
- [5] S.J.Brodsky and G.P.Lepage, Phys. Rev. D22 (1980) 2157.
- [6] L.Frankfurt, W.Koepf, and M.Strikman, Preprint TAUP-2290-95, hep-ph/9509311.
- [7] B. Blättel, G. Baym, L. L. Frankfurt and M. Strikman, Phys. Rev. Lett. 71 (1993) 896.
- [8] L. Frankfurt, G. A. Miller and M. Strikman, Phys. Lett. B304 (1993) 1.
- [9] S. J. Brodsky, L. Frankfurt, J. F. Gunion, A. H. Mueller and M. Strikman, Phys. Rev. D50 (1994) 3134.
- [10] F. E. Low, Phys.Rev. D12 (1975) 163.
- [11] J.Gunion and D.Soper, Phys.Rev.D15 (1977) 2617.
- [12] A.H.Mueller.Nucl.Phys.B335 (1990) 115
- [13] E.M. Levin, Sov.J.Nucl.Phys. 45(1987) 150
- [14] E. L. Feinberg and I. Y. Pomeranchuk, Suppl.Nuovo Cim. III (1956) 652.
- [15] M. L. Good and W. D. Walker, Phys.Rev.120, (1960) 1857.
- [16] H. Miettinen and J. Pumplin, Phys. Rev. D18, 1696 (1978);Phys. Rev. Lett. 42, 204 (1979).
- [17] H. Heiselberg, G. Baym, B. Blaettel, L.L. Frankfurt, M. Strikman, Phys. Rev. Lett. 67 (1991) 2946.
- [18] A.V. Radyushkin, Acta Physica Polonica, B12 (1995) 2067.
- [19] V. L. Chernyak and A. R. Zhitnitsky, Phys. Rep. 112 (1984) 173.
- [20] M.Gluck, E.Reya and V.Vogt, Z.Phys. C67,433 (1995)




FIG. 1. Leading small- x contribution to the forward virtual Compton amplitude.




FIG. 2. Comparison of $P_{\pi p}(\sigma)$ calculated in pQCD using eq.(6.14) and GRV parameterizations [20] of the gluon density and fits based on the analysis of the soft diffraction data [7].




FIG. 3. Incident momentum dependence of $P_{\pi p}(\sigma)$ for small σ calculated using eq. (6.14) and GRV NLO parameterization [20] of the gluon density.