

F_2^N at Low Q^2

Ioana Niculescu

The George Washington University

January, 2000

F_2^N at low Q^2

Submitted by

I. Niculescu (spokesperson)

The George Washington University, Washington, DC

C. S. Armstrong (spokesperson), R. Ent, K. Garrow, A. Lung, D. J. Mack,
J. Mitchell, S. A. Wood

Jefferson Lab, Newport News, VA

J. Arrington, D. De Schepper, D. Geesaman, B. Mueller, D. Potterveld, P. Reimer
Argonne National Laboratory, Argonne, IL

D. Meekins

Florida State University, Tallahassee, FL

O. K. Baker, H. Bitao, M. E. Christy, A. Cochran, P. Guèye, C. Jackson,
C. E. Keppel, Y. Liang, V. Nazarian, L. Tang

Hampton University, Hampton, VA

G. Niculescu

Ohio University, Athens, OH

P. Stoler

Rensselaer Polytechnic Institute, Troy, NY

E. Kinney

The University of Colorado at Boulder, Boulder, CO

D. B. Day


The University of Virginia, Charlottesville, VA

R. Asaturyan, H. Mkrtychyan, S. Stepanyan, V. Tadevosyan

Yerevan Physics Institute, Armenia

~30 collaborators
10 institutions

Inclusive electron scattering





$$x = \frac{Q^2}{2 M \nu}; \quad \xi = \frac{2x}{1+r}, \quad r = \sqrt{1 + \frac{4M^2 x^2}{Q^2}}$$

$$(Q^2 \rightarrow \infty \quad \xi \rightarrow x)$$


$$\frac{d^2\sigma}{d\Omega dE'} = \frac{4\alpha^2 E'^2}{Q^4} [2W_1(\nu, Q^2) \sin^2(\theta/2) + W_2(\nu, Q^2) \cos^2(\theta/2)]$$


$F_2 = \nu W_2$





Bloom-Gilman Duality






} SLAC
fit

} JLab
1998


} Stein
1978


10%
(R.C. + norm)


Scaling curve




Deuterium










Kinematics



— kinematic region of this experiment

Proposed Kinematics

E_b (GeV)	θ_e (deg.)	E' (GeV)	x	Q^2 (GeV/c) ²	ϵ
4.4	16.0	0.4-3.9 (22 settings)	0.016-1	0.13- <u>1.42</u>	0.18-0.95
4.4	10.5	0.4-3.9 (22 settings)	0.007-1	0.05-0.61	0.18-0.98
3.3	10.5	0.4-3.1 (20 settings)	0.007-1	0.04-0.36	0.24-0.98
2.2	10.5	0.4-2.2 (17 settings)	0.008-1	<u>0.03</u> -0.16	0.35-0.98

Take data on Hydrogen
Deuterium

Activity	Time (hours)
Data acquisition	68
Configuration changes	70
Beam energy changes	18
Checkout	12
Total	168 (7 days)

+ 1 (TAC)

Table 2: Beam time request for the proposed experiment.